

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/04, A61K 31/445	A1	(11) International Publication Number: WO 95/11894 (43) International Publication Date: 4 May 1995 (04.05.95)															
(21) International Application Number: PCT/US94/11790 (22) International Filing Date: 18 October 1994 (18.10.94) (30) Priority Data: 145,903 29 October 1993 (29.10.93) US (71) Applicants: THE UNIVERSITY OF TOLEDO [US/US]; 2801 West Bancroft Street, Toledo, OH 43606 (US). GLIATECH, INC. [US/US]; 23420 Commerce Park Road, Beachwood, OH 44122 (US). (72) Inventors: DURANT, Graham, J.; 55 Captain Luther Little Waye, Marshfield, MA 02050 (US). KHAN, Amin, M.; 6549 Park N, B-7, Solon, OH 44139 (US). TEDFORD, Clark, E.; 1448 Bell Road, S. Russell, OH 44022 (US). (74) Agents: LAWRENCE, Stanton, T. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published With international search report.															
(54) Title: HISTAMINE H ₃ -RECEPTOR ANTAGONISTS AND THERAPEUTIC USES THEREOF																	
<table border="1"> <caption>Food Intake (grams) Data</caption> <thead> <tr> <th>Treatment</th> <th>2 HR</th> <th>4 HR</th> </tr> </thead> <tbody> <tr> <td>VEHICLE</td> <td>~6.5</td> <td>~7.5</td> </tr> <tr> <td>2016 (3 mg/kg)</td> <td>~6.0</td> <td>~7.5</td> </tr> <tr> <td>2016 (10 mg/kg)</td> <td>~2.0</td> <td>~6.0</td> </tr> <tr> <td>2016 (30 mg/kg)</td> <td>~1.5</td> <td>~2.5</td> </tr> </tbody> </table>			Treatment	2 HR	4 HR	VEHICLE	~6.5	~7.5	2016 (3 mg/kg)	~6.0	~7.5	2016 (10 mg/kg)	~2.0	~6.0	2016 (30 mg/kg)	~1.5	~2.5
Treatment	2 HR	4 HR															
VEHICLE	~6.5	~7.5															
2016 (3 mg/kg)	~6.0	~7.5															
2016 (10 mg/kg)	~2.0	~6.0															
2016 (30 mg/kg)	~1.5	~2.5															
(57) Abstract <p>The present invention is directed to compounds of the class of piperidyl-imidazole derivative histamine H₃-receptor antagonist. Such compounds have affinity for histamine H₃-receptor, and preferably penetrate the blood-brain barrier. The compounds can block the soporific effect of an H₃-receptor agonist. Illustrative of the compounds of the invention is the molecule 4-(1-cyclohexylvaleroyl-4-piperidyl)-1H-imidazole. These compounds have been found to have utility in treating cognitive disorders and have been found to be useful as appetite suppressants.</p>																	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HISTAMINE H₃-RECEPTOR ANTAGONISTS
AND THERAPEUTIC USES THEREOF

The present application is a continuation-in-part of International application No. PCT/US93/03104, filed March 31, 1993 which is a continuation-in-part of application Serial No. 07/862,657, filed April 1, 1992, each of which is incorporated herein by reference in its entirety.

1. FIELD OF THE INVENTION

The present invention relates to novel compounds having potent activity as histamine H₃-receptor ("H₃") antagonists, and methods of using such compounds.


2. BACKGROUND OF THE INVENTION

Dementias tend to be characterized by cognitive disorders and often by depression. A particularly devastating dementia is Alzheimer's disease (AD). AD affects more than 30% of humans over 80 years of age, and as such, represents one of the most important health problems in developed countries (Evans et al., *J.A.M.A.* 262: 2551-2556 (1989); Katzman and Saitoh, *FASEB J.* 280: 278-286 (1991)). This neurodegenerative disorder of unknown etiology is clinically characterized by gradual impairment of cognitive function. The large buildup of intracytoplasmic neurofibrillary tangles and neurite plaques observed histopathologically in AD plausibly leads to degeneration of affected nerve cells. At least one study showed decreases in histamine and histidine levels in frontal, temporal and occipital cortices and in the caudate nucleus of brains from AD patients examined post mortem (Mazurkiewics and Wsonwah, *Can. J. Physiol. Pharmacol.*, 67:75-78 (1989)).

Histamine is a chemical messenger involved in various complex biological actions. It is widely distributed in the plant and animal kingdoms. In

SUBSTITUTE SHEET (RULE 26)

mammals, including man, it occurs mainly in an inactive bound form in most body tissues. When released, histamine interacts with specific
5 macromolecular receptors on the cell surface or within a target cell to elicit changes in many different bodily functions. Histamine (4(2-aminoethyl)imidazole) is a base. Its chemical structure is:



15 Histamine receptor pharmacology has revealed three subtypes of receptors which mediate (or are associated with) the activity of histamine. These receptors are most commonly referred to as H₁, H₂, and H₃. The most recently discovered of these receptors is the H₃
20 histamine receptor. Early studies suggested the presence of another histamine receptor when it was demonstrated that histamine inhibits its own synthesis and release in brain slices by a negative feedback process operating at the level of histaminergic nerve-
25 endings (see, for example, Arrang, J.M. *et al.* *Nature* 302:832-837 (1983)). More recently, the H₃ receptor has been shown to function as a pre-synaptic autoreceptor inhibiting histamine synthesis and histamine release from neurons, especially in the
30 control nervous system (Arrang, *et al.* *Nature* 327:117-123 (1987)). The presence of H₃ receptors in peripheral tissues has also been reported and here too they appear to be involved with the nervous system. Thus, histamine depresses sympathetic neurotrans-
35 mission in the guinea pig mesenteric artery by interacting with H₃ receptors on the perivascular nerve

SUBSTITUTE SHEET (RULE 26)

terminals (Ishikawa and Sperelakis, *Nature* 327:158 (1987)). This important observation suggests that histamine may control the release of other
5 neurotransmitters (Tamura et al., *Neuroscience* 25:171 (1988)). Inhibitory histamine H₂ receptors also exist in the guinea pig ileum where their role appears to be to modify the magnitude of histamine contraction, rather than affecting histamine release
10 (Trzeciakowski, *J. Pharmacol. Exp. Therapy* 243:847 (1987)). Particularly intriguing is the discovery of H₂ receptors in the lung (Arrang et al. *supra* (1987)). The presence of histamine H₂ receptors in the lung raises the question of whether they control histamine
15 release in anaphylaxis and whether they may be manipulated to provide therapy in asthma. Indeed it has been suggested that H₂ receptors may have a modulating role on excitatory neurotransmission in airways. Generally, however, H₂ receptor inhibition
20 tends to increase histamine activity, with potentially detrimental effects. Thus, it is desirable to avoid introducing H₂ receptor antagonists that act on peripheral tissues.

Histamine H₂ receptor activation was found to
25 inhibit acetylcholine release in a guinea pig ileum model (Poli et al., *Agents and Actions* 33: 167-169). Selective H₂-receptor blockers reversed the histamine-induced inhibitory effect. Histamine also decreased serotonin release; this effect was reversed with an
30 H₂-antagonist, and was suggested to operate via the histamine H₂-receptors (Schlicker et al., *Naunyn-Schmiedaberg's Arch. Pharmacol.* 337: 588-590 (1988). Activation of H₂-receptors was found to inhibit excitatory presynaptic potentials (Arrang et al., *J.*
35 *Neurochem.* 51:105 (1988)).

SUBSTITUTE SHEET (RULE 26)

One reported highly specific competitive antagonist of histamine H_2 receptors is thioperamide (Arrang et al., supra (1987)). Although thioperamide is a very potent antagonist *in vitro* ($K_i = 4.3$ nmol/L), relatively high doses are required *in vivo* to inhibit histamine release from the brain in rats (Ganellin et al., Collect. Czech. Chem. Commun. 56:2448-2455 (1991)). Ganellin et al. suggests that this most probably results from poor penetration through the blood-brain-barrier by this peramide, although the pharmacokinetic properties of thioperamide may also play a role. Moreover, the thiourea functionality found in thioperamide may result in higher intrinsic toxicity of thioperamide.

Thiourea-containing drugs are known to be associated with undesirable side effects in clinical use. For example, with thiourea-containing drug molecules that are used to treat hyperthyroidism, agranulocytosis is known to be a serious, and occasionally fatal, toxic effect in clinical use (see, e.g., Brimblecombe et al. Gastroenterology 74:339-346 (1978)). The thiourea-containing histamine H_2 -receptor antagonist metiamide caused a low incidence of granulocytopenia in peptic ulcer patients and was withdrawn from clinical use (Forrest et al., Lancet 1: 392-393 (1975)). In high dose, repeated dose toxicological studies in dogs, incidences of agranulocytosis were seen at 162 mg/kg/day (Brimblecombe et al., "Toxicology of Metiamide," International Symposium on Histamine H_2 - Receptor Antagonists, Wood and Simpkins, Smith Kline & French, pp. 53-72 (1973)). A proportion of dogs (<10%) died acutely with pulmonary edema and pleural effusion. The metiamide isostere cimetidine, in which the thiourea group was replaced by an alternative group

SUBSTITUTE SHEET (RULE 26)

(cyanoguanidine), did not cause granulocytopenia, or any other side effects in animal toxicity studies or in clinical usage by multimillions of patients, indicating that the toxicological problems with metiamide could be attributed to the presence of the thiourea group (Brimblecomb et al., *supra*). It is likely that the thiourea functionality, with its association with toxicological phenomena and its likelihood of inducing undesirable side effects, could limit the clinical development of thioperamide.

Although some predictions have been made concerning the ability of molecules to pass through the blood brain barrier, these predictions are at best speculative. The rate and extent of entry of a compound into the brain are generally considered to be determined primarily by partition coefficient, ionization constant(s) and molecular size. No single partition solvent system has emerged as a universally applicable model for brain penetration, although the octanol water system has received particular attention, and Hansch and coworkers have suggested that a partition coefficient in this system of about 100 is optimal for entry into the central nervous system (CNS) (Glave and Hansch, *J. Pharm. Sci.*, 61:589 (1972); Hansch et al., *J. Pharm. Sci.*, 76:663 (1987)). Comparisons between known H₂ antagonists, however, suggest that there is no such simple relationship between their brain penetration and octanol water partition coefficients (Young et al., *J. Med. Chem.* 31:656 (1988)). The comparison of the ability of histamine H₂ receptor antagonists to cross the blood brain barrier suggests that brain penetration may increase with decreasing over-all hydrogen binding ability of a compound (Young et al., *supra*). However, optimizing H₂ receptor antagonists to improve brain

SUBSTITUTE SHEET (RULE 26)

penetration reduced antagonist potency (Young et al.,
supra). Thus it is fundamentally difficult to
optimize both blood brain barrier permeability and
5 function of a compound.

It is therefore an object of the present
invention to provide novel potent histamine H₃-receptor
antagonists that are better able to penetrate the
blood-brain-barrier than previously reported
10 compounds.

Further it is an object of the present invention
to provide novel potent histamine H₃-receptor
antagonists that have reduced toxicity compared to
other known H₃ antagonists.

15 Another object of the present invention is to
provide histamine H₃-receptor antagonists that will act
selectively on the brain and have limited activity on
H₃ receptors in peripheral tissues.

It is yet another object of the present invention
20 to provide a novel class of histamine H₃-receptor
antagonists.


3. SUMMARY OF THE INVENTION

The present invention provides novel compounds
25 having activity as histamine H₃-receptor antagonists.
In a preferred aspect, the compounds of the invention
exhibit ready penetration of the blood-brain-barrier
and reduced toxicity. The novel compounds of this
invention include compounds of the formula:

30

35

SUBSTITUTE SHEET (RULE 26)



wherein D is CH_2 or CH_2-CH_2 , Z represents S or O, preferably O, x is 0 or 1, n is an integer from 0 to 6, R_1 represents preferably hydrogen, or a hydrolyzable group, but can be a lower alkyl or aryl group, and R_2 represents a linear chain, branched chain or carbocyclic group or aryl group of up to about 20 carbon atoms, and salts thereof. If R_2 is tert-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0. If R_2 is adamantane, the sum of x and n must be greater than 1. The various alkyl or aryl groups can have functional group substituents.

It has been discovered that amide or carbamate functional groups can be used to join alkyl or aryl substituents to the piperidyl nitrogen of 4(4-piperidyl)-1H-imidazole groups. Other cyclic imides, particularly pyrrolidyl or cycloheptamidyl ($C_6H_{11}N$) can be substituted for piperidine. In a preferred aspect, the compounds of the invention are surprisingly effective at transport across the blood brain barrier, thus limiting their effects primarily to cerebral histamine H_3 -receptors, and are also less toxic than histamine H_3 -receptor antagonists based on a thiourea functional group.

In addition, the present invention encompasses a pharmaceutical composition comprising a compound of

SUBSTITUTE SHEET (RULE 26)

the invention, and a method of using a compound or pharmaceutical composition of the invention in an animal, particularly in a human, to treat Alzheimer's disease and other dementias by ameliorating the cognitive defects and neurodegenerative effects associated therewith. The histamine H_3 -receptor antagonists of the invention have additional therapeutic uses where increased arousal and attention is desired.

Furthermore, the present invention encompasses a method of suppressing appetite in an animal, particularly a human, comprising administering to the animal an effective amount of a compound of the present invention. Thus, the invention encompasses the treatment of obesity in animals suffering from such a disorder. The invention further encompasses a pharmaceutical composition comprising a compound of the present invention; and optionally a pharmaceutically acceptable carrier. This pharmaceutical composition can be administered to suppress appetite in an animal, particularly a human.

4. BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Binding of N^{α} -methylhistamine to rat cortical homogenate. Open box: total bound; x'ed box: specific binding; closed box: non-specific binding.

FIG. 2. Binding of 3H -labeled N^{α} -methylhistamine to the cortical homogenate of thioperamide injected rats.

FIG. 3. Binding of 3H -labeled N^{α} -methylhistamine to the cortical homogenate of compound 1 injected rats.

FIG. 4. The effect of α -methylhistamine on sleeping one hour after injection.

SUBSTITUTE SHEET (RULE 26)

FIG. 5. The effect (dose-response) of thioperamide on sleep induced by R(-)- α -methylhistamine (30 mg/kg).


5 FIG. 6. The effect (dose-response) of compound 1 on sleep induced by R(-)- α -methylhistamine (25 mg/kg).

FIG. 7. The effect of compound 1 on 2 and 4 hour food intake in 24-hour-fasted rats. Solid bar: 2 hour food intake. Cross-hatched bar: 4 hour food intake.

10 FIG. 8. The effect of compound 1 on 6 hour food intake in 24-hour-fasted rats.

5. DETAILED DESCRIPTION OF THE INVENTION

15 The compounds of the present invention are compounds of the general formula.



wherein D is CH₂ or CH₂-CH₂, Z represents sulfur (S) or oxygen (O), preferably O, x is 0 or 1, n is an integer from 0 to 6, R₁ represents hydrogen, an *in vivo* hydrolyzable group, a lower alkyl group, a lower cyclic alkyl group, or a lower aryl group, and R₂ represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 carbon atoms including mono and bicyclic moieties, and a substituted or an

SUBSTITUTE SHEET (RULE 26)

unsubstituted aryl group of up to about 20 carbon atoms, or any combination of above-mentioned groups, or salts thereof. In a specific embodiment, R_2 can represent a disubstituted methyl, such as but not limited to dicyclohexyl methyl ($-\text{CH}(\text{C}_6\text{H}_{11})_2$), diphenyl methyl ($-\text{CH}(\text{C}_6\text{H}_5)_2$), and the like. If R_2 is tert-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0. If R_2 is adamantane, the sum of x and n must be greater than 1.

In a preferred embodiment, R_1 is hydrogen. It is also contemplated that R_1 can be a hydrolyzable leaving group, such as an acyl or carbamyl, including where $R_1 = -\text{CZ}(\text{O})_x(\text{CH}_2)_n\text{R}_2$, as in I above. It is well known that N-acylimidazoles are hydrolytically labile, and R_1 may be selected such that it yields the parent imidazole compound *in vivo* at an optimal rate. Such hydrolysis will yield the compound with hydrogen as R_1 . Thus, the contemplated compounds of the invention with a hydrolyzable substituent at R_1 are functionally equivalent to the preferred embodiment, i.e., where R_1 is hydrogen. R_1 can also be a lower linear chain, branched chain, or cyclic alkyl, or a lower aryl. The term "lower" as applied to the alkyl or aryl substituents at R_1 indicates the presence of up to seven carbon atoms. In specific embodiments *infra*, R_1 is methyl, benzyl, methylcyclohexane, N-cyclohexylformamide, benzaldehyde, and t-butylaldehyde.

In yet a further embodiment, the nitrogen atom at position 3 of the imidazole ring can be substituted with a lower alkyl or aryl group, or with a hydrolyzable leaving group.

In a preferred embodiment, D is $\text{CH}_2\text{-CH}_2$, resulting in a piperidine ring structure. However, it is contemplated that D can be CH_2 , yielding a pyrrolidine

SUBSTITUTE SHEET (RULE 26)

ring structure. In yet another embodiment, D can be $(CH_2)_3$, yielding a cycloheptimide (seven membered heterocycle with one nitrogen). While orientation of the imidazole group distal to the N of the piperidine is preferred, the invention contemplates the imidazole at the 2 or 3 position on the piperidine (or the 2 position of pyrrolidine, or the 2 and 3 position of the cycloheptimide ring). These alternate embodiments can be used instead of the piperidyl embodiment with the imidazole group located at the 4 position, although the piperidyl embodiment is preferred.

Although the present invention is not limited to any mechanistic theory, it is believed that the blood brain barrier is permeable to the compounds of the present invention in part because of the subtle decrease in polarity afforded by an amide or carbamate bond linking the $-(O)_x(CH_2)_nR$ moiety (e.g., a hydrophobic tail) to the 4(4-piperidyl)-1H-imidazole (or 4(3-pyrrolidyl)-1H-imidazole) structure. With slightly less polarity and hydrogen-bonding capability than urea or thiourea, the amide or carbamate functionality can more efficiently traverse the blood brain barrier. Moreover, the dipole of the amide or carbamate is distal to the hydrophobic tail, more proximal to the imidazole (which is a fairly polar group), and thus tends to effect greater amphiphilicity in the molecule. That the compounds of the invention retain amphiphilic character is important for solubility in aqueous solution. Solubility in aqueous solution is desirable for a compound to be used therapeutically in an animal particularly in a human. That such a subtle difference, use of an amide or carbamate functionality, should perceptibly alter blood brain barrier permeability may be considered to be surprising since it is not generally appreciated.

SUBSTITUTE SHEET (RULE 26)

In preferred embodiments, a bulky hydrocarbon R_2 group is chosen so that the net hydrophilicity of the H_3 -receptor antagonist is increased, and the steric effects of a bulky substituent at R_2 are decreased, by increasing the number of methylenes in a straight chain alkyl group (i.e., in Formula I, $n > 1$). In a specific embodiment, a tetramethylene bound to the amide or carbamate group is used. Preferably a cyclic alkyl or aryl group is linked to the amide or carbamate via the straight chain alkyl group. In a specific embodiment, tetramethylene cyclohexane (cyclohexylbutyl) is bound to an amide. Although specific hydrophobic alkyl and aryl groups have been mentioned, one of ordinary skill in the art will recognize that there are many possible hydrophobic groups for use in the compounds of the invention. These fall within the scope of the instant invention.

Thus, R_2 can be one or more bulky substituent groups. As stated above, in a preferred aspect of the invention, the bulky substituents are removed from the amide or carbanate group on the piperidyl-imidazole by increasing n . In one embodiment, R_2 is CHR_3R_4 , in which n is 3 or 4 and R_3 and R_4 are cyclohexyl, phenyl, or the like. R_3 and R_4 can be the same group or different groups. In another embodiment, R_2 is decalin or adamantane or the like. If R_2 is adamantane, preferably n is greater than 1, but the sum of x and n must be greater than 1.

As used herein, the phrase linear chain or branched chained alkyl groups of up to about 20 carbon atoms means any substituted or unsubstituted acyclic carbon-containing compounds, including alkanes, alkenes and alkynes. Examples of alkyl groups include lower alkyl, for example, methyl, ethyl, n -propyl, iso-propyl, n -butyl, iso-butyl or tert-butyl; upper

SUBSTITUTE SHEET (RULE 26)

alkyl, for example, octyl, nonyl, decyl, and the like; and lower alkylene, for example, ethylene, propylene, propyldiene, butylene, butyldiene, and the like. The
5 ordinary skilled artisan is familiar with numerous linear and branched alkyl groups, which are with the scope of the present invention.

In addition, such alkyl group may also contain various substituents in which one or more hydrogen
10 atoms has been replaced by a functional group. Functional groups include but are not limited to hydroxyl, amino, carboxyl, amide, ester, ether, and halogen (fluorine, chlorine, bromine and iodine), to mention but a few.

15 As used herein, substituted and unsubstituted carbocyclic groups of up to about 20 carbon atoms means cyclic carbon-containing compounds, including but not limited to cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, and the like. Such cyclic
20 groups may also contain various substituents in which one or more hydrogen atoms has been replaced by a functional group. Such functional groups include those described above, and lower alkyl groups as described above. The cyclic groups of the invention
25 may further comprise a heteroatom. For example, in a specific embodiment, R₂ is cyclohexanol.

As used herein, substituted and unsubstituted aryl groups means a hydrocarbon ring bearing a system of conjugated double bonds, usually comprising six or
30 more even number of π (pi) electrons. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, anisyl, tolyl, xylenyl and the like. According to the present invention, aryl also includes heteroaryl groups, e.g., pyrimidine or thiophene.
35 These aryl groups may also be substituted with any number of a variety of functional groups. In addition


SUBSTITUTE SHEET (RULE 26)

to the functional groups described above in connection with substituted alkyl groups and carbocyclic groups, functional groups on the aryl groups can be nitro groups.

As mentioned above, R_2 can also represent any combination of alkyl, carbocyclic or aryl groups, for example, 1-cyclohexylpropyl, benzyl cyclohexylmethyl, 2-cyclohexylpropyl, 2,2-methylcyclohexylpropyl, 2,2-methylphenylpropyl, 2,2-methylphenylbutyl.

In a specific embodiment, R_2 represents cyclohexane, and $n=4$ (cyclohexylvaleroyl). In another specific embodiment, R_2 represents cinnamoyl.

Particularly preferred are compounds of the formula:



wherein x is 0 or 1, n is an integer from 0 to 6, more preferably $n = 3-6$, and most preferably $n=4$, and R is as defined for R_2 above. Examples of preferred alkyl groups for R include but are not limited to cyclopentyl, cyclohexyl, adamantane methylene, dicyclohexyl methyl, decanyl and *t*-butyryl and the like. Examples of preferred aryl and substituted aryl groups include but are not limited to phenyl, aryl cyclohexyl methyl and the like.

SUBSTITUTE SHEET (RULE 26)

5.1. SYNTHESIS OF THE COMPOUNDS

The compounds of the present invention can be synthesized by many routes. It is well known in the art of organic synthesis that many different synthetic protocols can be used to prepare a given compound. Different routes can involve more or less expensive reagents, easier or more difficult separation or purification procedures, straightforward or cumbersome scale-up, and higher or lower yield. The skilled synthetic organic chemist knows well how to balance the competing characteristics of synthetic strategies. Thus the compounds of the present invention are not limited by the choice of synthetic strategy, and any synthetic strategy that yields the compounds described above can be used.

As shown in the Examples, *infra*, two general procedures can be used to prepare the instant compounds. Both involve condensation of an activated (electrophilic) carbonyl with the nucleophilic piperidyl nitrogen of 4-(4-piperidyl)-1H-imidazole.

The first procedure involves preparing the acid chloride derivative or acid anhydride of a carbonyl, i.e., activating the carbonyl. This activated carbonyl is added in molar excess to the piperidyl-imidazole in the presence of a molar excess of an unreactive base, for example, but not limited to, dicyclohexyl amine.

The second procedure is to condense the piperidyl-imidazole with a slight molar excess of a dicarbonate, again in the presence of an unreactive base, for example and not by way of limitation, triethylamine. This method can be used especially in the preparation of carbamate compounds.

A preferred synthesis of the 4-(4-piperidyl)-1H-imidazole is also provided. Commercially available 4-

SUBSTITUTE SHEET (RULE 26)

acetyl pyridine (Aldrich Chemical Co.) is converted into the key intermediate 4-(4-pyridyl)-1H-imidazole by bromination with hydrogen bromide in acetic acid
5 (Barlin, et al., Aust. J. Chem. 42:735 (1989)) to yield the bromoacetyl pyridine in high yield. Reaction of bromoacetyl pyridine with formamide at 110°C affords the substituted imidazole in high yield. The reaction is usually performed without the addition
10 of any solvent. The pyridyl moiety is reduced by catalytic hydrogenation using 5-10% Rhodium on carbon in acidified water at a pressure of 20-55 atmospheres to yield 4-(4-piperidyl)-1H-imidazole. This synthesis is disclosed more fully in copending United States
15 patent application Serial No. 07/862,658, filed by the instant inventors on April 1, 1992, entitled "PROCESS FOR THE PREPARATION OF INTERMEDIATES USEFUL FOR THE SYNTHESIS OF HISTAMINE RECEPTOR ANTAGONISTS," which is specifically incorporated herein by reference in its
20 entirety.

Solvents for use in the synthesis of the compounds of the invention are well known in the art. The solvent must be non-reactive, and the starting materials and base must be soluble in the solvent.
25 Preferably, an aprotic organic solvent of medium to high polarity is used. For example, acetonitrile, can be used. Under appropriate conditions, in the synthesis of carbamates of the invention, an alcohol, e.g., methanol, can be used.

30 The electrophilic carbonyl group, which contains the R₂ moiety, can be obtained from commercial sources, or it may be prepared synthetically. In specific examples, *infra*, the carbonyl is obtained commercially. Activation of carbonyls is well known.
35 The acid chloride can be prepared by reacting the carboxylic acid with sulfonyl chloride.

SUBSTITUTE SHEET (RULE 26)

Alternatively, the acid chloride may be available commercially. In specific embodiments, *infra*, acid chlorides were obtained from commercial sources (Aldrich Chemical). Similarly, the acid anhydride can be prepared conveniently by reaction of a salt of the carboxylic acid with the acid chloride. In a specific embodiment, the carboxylic acid is reacted with a carbonate acid chloride to form an asymmetric acid anhydride. In another embodiment, the acid anhydride can be obtained commercially. In a specific embodiment, *infra*, the acid anhydride was obtained from Aldrich Chemical. Dicarbonates for use in the invention are available commercially, e.g., from Aldrich Chemical.

5.2. BIOLOGICAL ACTIVITY

The compounds of the present invention are biologically active in assays for histamine H₃-receptor antagonist activity, as well as in a radioligand binding assay in rat brain membranes (e.g., Table I, *infra*). The binding assay procedure used and its standardization with known H₃-receptor antagonists is shown in the examples *infra*.

Further biological studies can demonstrate that the histamine H₃-receptor antagonists of this invention reverse the soporific effects of the histamine H₃-receptor agonist, R(-)-alphamethylhistamine in mice when both drugs are administered peripherally (*infra*). In a specific embodiment, the compound designated No. 2016 reverses the soporific effect of R(-)-alphamethylhistamine.

The data in the Examples, *infra*, support the view that antagonists of histamine H₃-receptors of the invention are useful regulators of the sleep-

SUBSTITUTE SHEET (RULE 26)

wakefulness cycle with potentially useful cognitive and behavioral effects in mammals including humans.

In vivo studies can be used to show effectiveness of a compound of the invention to cross the blood-brain barrier, as shown in the examples, *infra*. The data support the view that drugs of the present invention penetrate the blood brain barrier and are able to exert beneficial central actions in mammals when these drugs are administered to the peripheral circulation.

5.3. THERAPY

The histamine H₃-receptor antagonists of the invention can be provided therapeutically for the treatment of a subject suffering from a cognitive disorder or an attention or arousal deficit, according to the present invention. One of ordinary skill in the art would readily determine a therapeutically effective dose of an H₃ receptor antagonist of the invention based on routine pharmacological testing and standard dosage testing. In one aspect of the present invention, the compounds can be administered in doses of about 0.01 to about 200 mg/kg, more preferably 1 to 100 mg/kg, and even more preferably 30 to 100 mg/kg. In a specific embodiment, greater than about 20 mg/kg of a compound of the invention was effective to reduce the soporific effect of (R) α -methylhistamine. Included in the routine pharmacological testing are toxicity studies to determine an upper limit dose. Such toxicity studies can include LD₅₀ studies in mice, and 15 day toxicity studies in mammals.

The histamine H₃-receptor antagonists of the invention are believed to increase the release of cerebral histamine, acetylcholine and serotonin. These compounds can lead to increased arousal and

SUBSTITUTE SHEET (RULE 26)

attention. They can also be of benefit in the treatment of cognitive disorders.

Therapy with a compound of the invention is indicated to treat dementia, as either a primary or an adjunct therapy. The compounds of the invention have clinical utility in the treatment of dementia disorders in general. In a preferred embodiment, a compound of the invention can be used in the treatment for Alzheimer's disease. The compounds can also be used to treat presenile and senile dementia, Huntington's chorea, tardive dyskinesia, hyperkinesia, mania, Tourette syndrome and Parkinson's disease, to name but a few. Other specific indications include the treatment of narcolepsy and hyperactivity in children. In another embodiment, the compounds of the invention can be used in the treatment of certain psychoses, for example forms of depression or schizophrenia.

The compounds of the invention can be used to arouse victims of comas induced by stroke, drugs or alcohol. In another embodiment, the compounds of the invention can be used to increase wakefulness, where this effect is desired. For example, the compounds of the invention, which are preferentially targeted to H₂ receptors in the brain, can be used to counteract the soporific effect of some antihistamines without negating the therapeutic effects of the antihistamines on peripheral tissue, e.g., lung. Thus allergy patients can relieve some of the side effects of antihistamine therapy. Similarly, the compounds of the invention can be used to reverse overdose of barbiturates and other drugs.

35

SUBSTITUTE SHEET (RULE 26)

5.3.1. Appetite Suppression

The histamine H₃-receptor antagonists of the present invention have been found to be useful in
5 animals, particularly humans, as appetite suppressants. Thus, in another embodiment of the present invention, these compounds are used to control weight gain, to treat obesity and to promote weight loss, as well as any other condition wherein appetite
10 suppression is desirable.

The present invention provides a method of suppressing appetite in an animal, including but not limited to a human, which comprises administering to the animal, an effective amount of a compound of the
15 present invention.

In specific embodiments, the present invention encompasses a method of suppressing appetite in an animal, including but not limited to a human, which comprises administering to an animal, including a
20 human, an effective amount of a compound of formula I or formula II, or a pharmaceutically acceptable salt thereof.

In a most preferred embodiment, the present invention encompasses a method for suppressing
25 appetite in an animal, including but not limited to a human, which comprises administering to an animal, including a human, an effective amount of 4-(1-cyclohexylvaleroyl-4-piperidyl)-1H-imidazole (compound 1) or a pharmaceutically acceptable salt thereof.

30 The invention also provides pharmaceutical compositions useful as appetite suppressants which comprise an effective amount of a compound of the present invention or a pharmaceutically acceptable salt thereof; and optionally a pharmaceutically
35 acceptable carrier or excipient. In particular, the present invention encompasses a pharmaceutical

SUBSTITUTE SHEET (RULE 26)

composition comprising an effective amount of 4-(1-cyclohexylvaleroyl-4-piperidyl)-1H-imidazole (compound 1) or a pharmaceutically acceptable salt thereof; and
5 optionally a pharmaceutically acceptable carrier or excipient.

An effective amount is an amount sufficient to achieve appetite suppression. One of ordinary skill in the art can readily determine an effective amount
10 of a compound of the present invention based on routine pharmacological testing and standard dosage testing. In one embodiment of the present invention, an effective amount of the compounds disclosed here is about 0.01 to about 200 mg/kg, more preferably 0.1 to
15 100 mg/kg, and even more preferably 30 to 100 mg/kg. The doses of the present invention can be given in divided or multiple doses over time as needed by the particular subject.

It is noted that the terms "appetite suppression" or "suppressing appetite" are known to those skilled
20 in the art and these are used herein consistently therewith. These terms as used herein include a reduction, decrease or amelioration in appetite, a reduction, decrease or amelioration in the desire or
25 craving for food, a reduction, decrease or amelioration in food intake. Appetite suppression can result in weight loss or weight control as desired. In a specific embodiment, the appetite suppressant of the present invention can be used to treat obesity and
30 any related condition in which weight control is desired, including but not limited to severe and moderate obesity as well as overweight subjects.

5.4. PHARMACEUTICAL COMPOSITIONS AND METHODS OF ADMINISTRATION

35

The effective dose of a compound of the invention, and the appropriate treatment regime can

SUBSTITUTE SHEET (RULE 26)

vary with the indication and patient condition, e.g., the treatment of a dementia or the treatment of tiredness may require different doses and regimens.

- 5 These parameters are readily addressed by one of ordinary skill in the art and can be determined by routine experimentation.

- A therapeutic compound of the invention may also contain an appropriate pharmaceutically acceptable
10 carrier or excipient, diluent or adjuvant, i.e., the compound can be prepared as a pharmaceutical composition. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic
15 origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid
20 carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium carbonate, magnesium stearate, sodium stearate, glycerol monostearate,
25 talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. These compositions can take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained-release formulations and the like. When
30 administering the compounds of the present invention as an appetite suppressant, the preferred dosage form is orally by tablet, pill or capsule. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such
35 compositions will contain an effective therapeutic amount of the active compound together with a suitable

SUBSTITUTE SHEET (RULE 26)

amount of carrier so as to provide the form for proper administration to the patient. While intravenous injection is a very effective form of administration, other modes can be employed, including but not limited to intraventricular, intramuscular, intraperitoneal, intra-arteriolar, and subcutaneous injection, and oral, transdermal, nasal and parenteral administration.

10 The therapeutic agents of the instant invention may be used for the treatment of animals, and more preferably, mammals, including humans, as well as mammals such as dogs, cats, horses, cows, pigs, guinea pigs, mice and rats.

15 In another embodiment, the therapeutic compound can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

In yet another embodiment, the therapeutic compound can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann.*

SUBSTITUTE SHEET (RULE 26)

Neurol. 25:351 (1989); Howard et al., *J. Neurosurg.* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, supra, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

6. EXAMPLES

A series of compounds were prepared and tested for their histamine H₃ receptor antagonist activity. The results are summarized in Table 1. The antagonist activity of the compounds was detected by observing inhibition of (³H)-N-(alpha)methylhistamine activity on rat brain membranes.


6.1. SYNTHESIS OF THE COMPOUNDS

The amide and carbamate compounds of Table 1 were synthesized from 4-(4-piperidyl)-1H-imidazole by three general procedures:

Procedure A: 4-(4-piperidyl)-1H-imidazole and the appropriate acid chloride were conjugated using dicyclohexylamine as base according to the following scheme:

SUBSTITUTE SHEET (RULE 26)

Scheme I




10


Procedure B: 4-(4-piperidyl)-1H-imidazole and the corresponding acid anhydride were conjugated using triethylamine as base according to the following scheme:

15

Scheme II



20



25

Procedure C: 4-(4-piperidyl)-1H-imidazole and the corresponding dicarbonate were conjugated using triethylamine as a base according to the following scheme:

30


6.1.1. PREPARATION OF 4-(1-CYCLOHEXYLVALEROYL-4-PIPERIDYL) 1H-IMIDAZOLE (COMPOUND 1)

35

To a mixture of 755 mg (5.00 mmol) 4-(4-piperidyl)-1H-imidazole and 942 mg (5.20 mmol) of

SUBSTITUTE SHEET (RULE 26)

Scheme III



10 dicyclohexylamine in 10 ml anhydrous acetonitrile at 25°C was slowly added 1.06 g (5.20 mmol) cyclohexanevaleroyl chloride in 2 ml of dichloromethane over a period of 10 min with stirring; then the reaction mixture was heated at 60°C for 1.5 h. After

15 cooling to ambient temperature, the solid side product that was obtained (dicyclohexylammonium chloride) was filtered off and the filtrate was concentrated in vacuo to remove acetonitrile. The resulting crude oil was crystallized with methanol: anhydrous diethyl

20 ether to give 1.085 mg of analytically pure product as a yellow powder. Yield: 68%; M.P.: 159°C; MS: m/e=317(M⁺); ¹H NMR (CDCl₃): imidazole H: δ 7.65 (s, 1H), 6.75 (s, 1H); cyclohexylbutyl: δ 2.20 (m, 8H), 1.20 (m, 11H); piperidyl: 4.65 (d, 2H), 3.95 (d, 2H),

25 3.10 (d, 2H), 2.84 (m, 1H), 2.20 (m, 2H).

Compounds No. 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12 in Table I were synthesized in similar manner, i.e., by condensation of the acid chloride with 4(4-piperidyl) 1-H-imidazole in the presence of

30 dicyclohexylcarbodiimide. Purified product was obtained by preparative TLC Silica Gel GF. 60 (2000 Microns) and the solvent of recrystallization was methanol:anhydrous ether (20:80).

Compound No. 3, yield: 70%; oil; MS m/e 275 (M⁺);

35 ¹H NMR (CDCl₃): imidazole H: δ 7.60 and 6.75 (s, 1H); piperidine H: complex, δ 4.65 (d, 2H), 3.90 (d, 2H),

SUBSTITUTE SHEET (RULE 26)

3.10 (m, 3H), 2.10 (m, 2H); cyclohexyl acetyl H: δ 1.50 (m, 11H), 2.80 (m, 2H).

Compound No. 4, yield: 67%; oil; MS: m/e 267

5 (M+); ^1H NMR (CDCl_3): imidazole H: δ 7.50 and 6.60 (s, 1H); piperidine H: complex, δ 3.90 (d, 2H), 2.80 (m, 3H), 2.55 (m, 2H), 1.80 (m, 2H); phenyl acetyl H: δ 7.10 (m, 5H), 1.50 (m, 2H).

Compound No. 5, yield: 71%; oil; MS: m/e 297

10 (M+); ^1H NMR (CDCl_3): imidazole H: δ 7.80 and 6.70 (s, 1H); piperidine H: complex, δ 4.60 (d, 2H), 3.80 (d, 2H), 3.10 (m, 3H), 1.80 (d, 2H); phenyl propyl H: δ 7.20 (m, 5H), 2.65 (m, 2H), 2.35 (m, 2H), 2.10 (m, 2H).

Compound No. 6, yield: 74%; oil; MS: m/e 289

15 (M+); ^1H NMR (CDCl_3): imidazole H: δ 7.70 and 6.80 (s, 1H); piperidine H: complex, δ 4.60 (d, 2H), 3.85 (d, 2H), 3.10 (m, 3H), 1.90 (m, 2H); cyclohexyl ethyl H: δ 1.10 (m, 11H), 2.00 (br, 2H), 2.20 (m, 2H).

Compound No. 7, yield: 75%; oil; MS: m/e 283

20 (M+); ^1H NMR (CDCl_3): imidazole H: δ 7.60 and 6.70 (s, 1H); piperidine H: complex, δ 4.60 (d, 2H), 3.90 (d, 2H), 3.10 (m, 3H), 1.80 (m, 2H); phenyl ethyl H: δ 7.30 (m, 5H), 2.10 (br, 2H), 1.50 (m, 2H).

Compound No. 8, yield: 69%; M.P.: 151°C; MS: m/e

25 327 (M+); ^1H NMR (CDCl_3): imidazole H: δ 7.65 and 6.80 (s, 1H); piperidine H: complex, δ 4.70 (d, 2H), 4.50 (d, 2H), 3.60 (m, 1H), 2.80 (m, 2H), 2.10 (m, 2H); adamantyl acetyl H: δ 1.80 (m, 12H), 3.10 (m, 2H), 4.05 (m, 1H).

Compound No. 9, yield: 62%; M.P.: 148°C

30 (decomposed); MS: m/e 357 (M+); ^1H NMR (CDCl_3): imidazole H: δ 7.60 and 6.85 (s, 1H); piperidine H: complex, δ 4.50 (d, 2H), 4.05 (m, 3H), 3.40 (d, 2H), 2.10 (m, 2H); dicyclohexyl acetyl H: δ 1.50 (m, 22H),
35 2.50 (m, 1H).

SUBSTITUTE SHEET (RULE 26)

Compound No. 10, yield: 64%; oil; MS: m/e 281 (M+); ¹H NMR (CDCl₃): imidazole H: δ 7.75 and 6.60 (s, 1H); piperidine H: complex, δ 4.70 (d, 2H), 4.20 (m, 3H), 2.80 (m, 2H), 2.10 (d, 2H); phenyl vinyl H: δ 7.40 (m, 5H), 6.50 (m, 2H).

Compound No. 11, yield: 62%; oil; MS m/e 351 (M+); ¹H NMR (CDCl₃): imidazole H: δ 7.50 and 6.40 (s, 1H); piperidine H: complex, δ 4.60 (d, 2H) 4.10 (m, 3H), 2.80 (d, 2H), 1.80 (m, 2H); phenyl cyclohexyl acetyl H: δ 7.20 (m, 5H), 1.80 (m, 11H), 3.70 (m, 1H).

Compound No. 12, yield: 72%; M.P.: 136°C; MS:m/e 304 (M+); ¹H NMR (CDCl₃): imidazole H: δ 7.70 and 6.80 (s, 1H); piperidine H: complex, δ 4.60 (d, 2H), 4.00 (m, 2H), 3.60 (m, 3H), 1.88 (m, 2H); cyclohexyl propyl H; complex, δ 1.20 (m, 17H).

6.1.2. ALTERNATIVE METHOD FOR THE PREPARATION OF 4-(1-CYCLOHEXYLVALEROYL-4-PIPERIDYL) 1H-IMIDAZOLE (COMPOUND 1)

Preparation of acid anhydride: Triethylamine (1.01 g, 10.00 mmol) was slowly added to a stirred solution of 1.84 g (10.00 mmol) cyclohexylpentanoic acid in 60 ml acetonitrile at 0°C. After 30 min. of stirring, 1.08 g (10.00 mmol) of ethylchloroformate was added slowly in 5-7 min., so that the temperature remained between 0°C and 5°C. After 1h stirring, the solution was used for the preparation of Compound 1.

Preparation of Compound 1: The freshly prepared acid anhydride was poured into a suspension of 1.54 g (10.20 mmol) of 4(4-piperidyl)imidazole and 1.03 g (10.20 mmol) triethylamine in 70 ml acetonitrile. After 1 h of heating at 80°C, the solution was concentrated under reduced pressure, and the oily residue was taken up with 75 ml water and then extracted with 150 ml ethylacetate. The residual oil

SUBSTITUTE SHEET (RULE 26)

was obtained, which crystallized on addition of ethylacetate/hexane. Yield: 74%.

This method provides the desired amide in good yield when the piperidylimidazole is added in slight molar excess, e.g., about a 1.01 to 1 molar ratio, to the asymmetric anhydride.

Compounds No. 52-58 in Table I were synthesized in similar manner, i.e., by condensation of the asymmetric ethylchloroformate acid anhydride with 4(4-piperidyl) 1H-imidazole in the presence of triethyl amine.

Commercially available 3,3-diphenylpropionic acid and 4,4-diphenylbut-3-enoic acid were used as the starting materials for compounds 52 and 54, respectively. The unsaturated alkene bond of 4,4-diphenylbut-3-enoic acid was reduced under mild conditions by Pd/C (5%)/H₂ catalysis. This intermediate was then used to synthesize compound 53. Both intermediates 3,3-diclohexylpropionic acid and 4,4-dicyclohexylbutanoic acid, used in the preparation of compounds 55 and 56, respectively, were prepared by reduction of 3,3-diphenylpropionic acid and 4,4-diphenylbutanoic acid in the presence of catalyse Rh/alumina (5%)/H₂, 5 atm.

Compound No. 52, yield: 69%; MS.:M/e 359 (M+); ¹H NMR CDCl₃:imidazole H: δ 7.50 and 6.70 (s, 1H); piperidine H: complex δ 4.60 (d, 2H), 3.10 (m, 3H), 2.60 (d, 2H), 1.40 (m, 2H); propionyl H: complex δ 3.05 (m, 1H), 2.00 (d, 2H); biphenyl H: complex δ 7.20 (m, 10H), MA.: calc. C=76.85, H=7.00, N=11.68; found, 76.32, 6.72, 10.89, respectively.

Compound No. 53, yield: 73%; MS.:M/e 373 (M+); ¹H NMR CDCl₃:imidazole H: δ 7.65 and 6.70 (s, 1H); piperidine H: complex δ 4.60 (d, 2H), 3.00 (m, 2H), 2.50 (m, 2H), 1.80 (m, 2H); butanoyl H: δ 3.05 (m,

SUBSTITUTE SHEET (RULE 26)

2H), 2.40 (m, 2H), 3.40 (m, 2H); diphenyl H: δ 7.10 (m, 10H).

Compound No. 54, yield: 64%; MS.: M/e 371 (M+); ¹H NMR CDCl₃: imidazole H: δ 7.40 and 6.50 (s, 1H);
5 piperidyl H: complex δ 4.50 (d, 2H), 3.60 (m, 3H), 3.20 (d, 2H), 1.50 (d, 2H); butenyl H: complex δ 6.70 (d, 1H), 3.50 (d, 2H); diphenyl H: δ 7.10 (m, 10H).

Compound No. 55, yield: 75%; MS.: M/e 371 (M+);
10 ¹H NMR CDCl₃: imidazole H: δ 8.00 and 7.10 (s, 1H); piperidyl H: complex δ 4.50 (d, 2H), 3.10 (d, 2H), 2.80 (m, 3H), 1.90 (d, 2H); propionyl H: complex δ 2.60 (d, 2H), 2.00 (m, 1H); dicyclohexyl H: complex δ 1.50 (m, 22H).

Compound No. 56, yield: 68%; MS.: M/e 385 (M+);
15 ¹H NMR CDCl₃: imidazole H: δ 8.00 and 7.05 (s, 1H); piperidyl H: complex δ 4.50 (d, 2H), 3.80 (d, 2H), 3.00 (m, 3H), 2.10 (m, 2H); butanoyl H: δ complex 2.80 (m, 2H), 1.80 (m, 2H), 1.40 (m, 1H); dicyclohexyl H:
20 complex δ 1.20 (m, 22H).

6.1.3. PREPARATION OF 4-(t-BUTOXY CARBONYL-4-PIPERIDYL) 1H-IMIDAZOLE (COMPOUND 2)

To a suspension of 224 mg (1.00 mmol) of 4-(4-piperidyl)-1H-imidazole dihydrochloride in 10 ml of
25 methanol was added 202 mg (2.00 mmol) of triethylamine (the suspension turned to a clear solution) followed by dropwise addition of 218 mg (1.00 mmol) of di-t-butyl dicarbonate in 5 ml methanol over a period of 10
30 min. The reaction mixture was stirred at 25°C for 6 h, at the end of which the volatile materials were removed in vacuo. The oily residue was partitioned between 50 ml chloroform and 25 ml water. The organic layer was washed with 50 ml brine solution, then dried
35 over anhydrous sodium sulfate. After filtration and removal of solvent, a pale yellow oil was obtained. The oil was treated with a mixture of methanol:

SUBSTITUTE SHEET (RULE 26)

petroleum ether (10:90). The resulting mixture was agitated vigorously with a glass rod until a solid appeared. After filtration and drying, the desired product was obtained as a white power. Yield: 65%; M.P.: 198°C; MS: m/e 251 (M⁺); ¹H NMR (CDCl₃): imidazole H: δ 7.60 (s, 1H) and 6.60 (s, 1H); piperidine H: δ 4.20 (d, 2H), 2.80 (m, 4H), 2.20 (d, 2H), 1.60 (m, 1H), t-BOC H: 1.45 (s, 9H).

Compounds No. 13 and 14 in Table I were synthesized in similar manner. The pure product was obtained by preparative TCL Silica GEL GF, 60 (2000 microns), and the solvent of recrystallization was methanol:anhydrous ether (20:80).

Compound No. 13, yield: 78%; M.P.: 180°C; MS: m/e 255 (M⁺); ¹H NMR (DMSO-d₆): imidazole H: δ 7.95 and 6.80 (s, 1H), NH: δ 7.80 and 6.60 (d, 1H); piperidine H: complex, δ 4.50 (d, 2H), 3.60 (m, 3H), 3.10 (m, 1H), 2.75 (m, 2H); phenyl H: δ 7.40 (m, 5H); MA: (C,H,N): 70.36%, 6.71%, 16.30%.

Compound No. 14, yield: 72%; M.P.: 185°C; ¹H NMR (CDCl₃): imidazole H: δ 7.60 and 6.80 (s, 1H); piperidine H: complex, δ 4.50 (d, 2H), 3.00 (m, 3H), 2.05 (d, 2H), 1.60 (m, 2H); t-butyl H: δ 1.10 (s, 9H).

6.1.4. PREPARATION OF 4(-4-PIPERIDYL)-1H-IMIDAZOLE

In a preferred embodiment, 4(4-piperidyl)-1H-imidazole for use in the synthesis of the H₃-receptors antagonists is prepared according to the following method.

Bromination of 4-acetyl piperidine (Aldrich) in hydrogen bromide/acetic acid was performed as described (Barlin *et al.*, Aust. J. Chem 42:735 (1989)).

SUBSTITUTE SHEET (RULE 26)

A mixture of 11.23g (4.00 mmol) of bromoacetyl pyridine and 3.98 ml (10.0 mmol) formamide were fused together at 110°C with stirring for 4h. The crude
5 reaction mixture was then concentrated on the rotary evaporator to remove volatile matter. The residue was dissolved in 50 ml methanol, and to this solution was added 100 ml anhydrous dimethyl ether slowly with stirring, which led to the formation of a brown
10 precipitate. After stirring for another 0.5h, the precipitate was filtered, washed with 50 ml anhydrous ether and dried. This solid residue was dissolved in 20 ml water and the aqueous solution was basified to pH 9 with sodium carbonate. To this solution was
15 added 150 ml absolute ethanol slowly with stirring till a solid formed, which was filtered off. The filtrate was heated to boiling, then treated with activated carbon and filtered. The filtrate was concentrated on rotary evaporator to dryness. Yield:
20 3.36g 58%; M.P.: 152°C (decomposed); MS: m/e 145 (M+), ¹H NMR (D₂O): imidazole H: δ 7.80 (s, 1H) and 7.20 (s, 1H); pyridyl H: 8.10 (d, 2H), 7.17 (d, 2H). The pyridyl moiety was reduced by catalytic hydrogenation using 5-10% rhodium on carbon in acidified water at
25 20-55 atmospheres as described (Schunack, Archiv. Pharma. 306:934 (1973)).

6.2. ANTAGONIST ACTIVITY IN VITRO

The various compounds were tested for the ability
30 to bind to the histamine H₂ receptor. A binding assay in a rat brain membrane preparation, based on inhibition of binding of [³H]-N-alpha-methylhistamine using excess unlabeled alpha-methylhistamine to account for nonspecific binding, was developed.
35 Total, specific and nonspecific binding of [³H]-N-alpha-methylhistamine to brain membranes is shown in

SUBSTITUTE SHEET (RULE 26)

FIG. 1. The K_d value was 0.19 nM in this preparation and the nonspecific binding was less than 20% of the total binding at the K_d value. The compounds

5 thioperamide (Arrang et al., Nature 327:117-123 (1987)) and burimamide (Black et al., Nature 236:385-390 (1972)) were tested as controls for this assay. The results are shown in Table I.

10

15

20

25

30

35

SUBSTITUTE SHEET (RULE 26)








- 34 -

TABLE I

4-Piperidyl (imidazole) Compounds and
Their Activities on Rat Brain Membranes.

5

(³H-N^α-methylhistamine as Radioligand)







Cmpd No.	R ₁	X (= CO-(O) _n (CH ₂) _n R)	IC ₅₀ (nm)	M.P.
Thioparamide	H		4.0±0.6 n=4	170°C
		Burimamide	156±57	
1	H		23±6 n=3	159°
3	H		19±12 n=3	Oil
4	H		1400±437 n=3	Oil
5	H		262±9 N=3	Oil
6	H		34±1.4 n=3	Oil
7	H		34.1±3.6 n=3	Oil
12	H		41.4±9 n=3	136°C

SUBSTITUTE SHEET (RULE 26)

	Cmpd No.	R ₁	X (= CO-(O) _n (CH ₂) _n R)	IC ₅₀ (nM)	M.P.
5	13	H		151 ± 44 n=4	180°C
	41	H		inactive n=2(1 μM)	192°C
10	42	CH ₃		inactive n=2(1 μM)	Oil
15	43	X		inactive n=3	99°C
	44	X		inactive n=2	81°C
20	45	X		inactive n=2	79°C
	46	PhCH ₂		inactive n=2	62°C
25	47	H		231 n=1	185°C
30	48	H		inactive n=2	168°C
35	52	H		93.1	129- 131°C









SUBSTITUTE SHEET (RULE 26)

- 36 -

Cmpd No.	R ₁	X (= CO-(O) _n (CH ₂) _n R)	IC ₅₀ (nm)	M.P.
53	H		124	oil
54	H		1000	158°C decomp.
55	H			118°C decomp.
56	H			152°C decomp.
57	H			
58	H			

SUBSTITUTE SHEET (RULE 26)

- 37 -

Compd. No.	Structure	IC ₅₀ (Nm)	M.P.
50		inactive n=2(μM)	148.5°-150.5°C
2		243.5±1.9 n=2	198°C
14		inactive n=2	185°C
8		inactive n=2	151°C
9		inactive n=2	148°C
10		570±172 n=3	Oil
11		260±38 n=2	Oil
51			115°C

SUBSTITUTE SHEET (RULE 26)

6.3. DISCUSSION

The results in Table I show that the compounds of the invention are effective for binding to the histamine H_3 -receptor. Interestingly, cyanoguanidine derivatives (e.g., compounds 47, 48 and 50) were ineffective at binding to the H_3 -receptor. This result is in contrast to earlier observations about H_2 -receptor antagonists. With H_2 -receptor antagonists, cyanoguanidine and thiourea-containing derivatives (cimetidine and metiamide, respectively) were found to be bioisosteres, i.e., functionally substantially equivalent (Brimblecombe et al., *Gastroenterology* 74: 339-347 (1978)).

7. PHARMACOLOGICAL EVALUATION IN THE CNS

A representative compound, 1, was tested in vivo for (1) the ability to penetrate the blood brain barrier; and (2) the effect of behavior in mice.

7.1. PENETRATION OF THE BLOOD-BRAIN BARRIER

Blood-brain barrier penetration in rats was assessed by an ex vivo binding procedure. Young adult male Long-Evans rats were injected i.p. with saline or H_3 antagonists in saline. At various times after injection animals were sacrificed, the cortex was removed, homogenized in 50 mM Na/K-phosphate buffer, pH 7.4, and the binding of 1 nM [3H]- N^α -methylhistamine was measured using 400 μ g protein of the homogenate. Nonspecific binding was accounted for by the inclusion of excess thioperamide in some samples. Under these conditions, the binding was approximately 90% specific.

As shown in FIG. 2, thioperamide at doses of 2, 5, and 10 mg/kg, when measured 15 min after injection, decreased the binding of [3H]- N^α -methylhistamine to H_3

SUBSTITUTE SHEET (RULE 26)

receptors in the cortex. This means that the thioperamide at these doses and after this time was able to penetrate the blood-brain barrier. Figure 3 shows that compound 1 also penetrates the blood-brain barrier one hour after injections of doses of 50 and 70 mg/kg. Taking into account the difference in affinity comparing thioperamide (4.0 nM) and compound 1 (23 nM), these data suggest that compound 1 penetrates the blood-brain barrier at least as well as thioperamide.

7.2. BEHAVIORAL EFFECTS IN MICE

The overall strategy to show central nervous system antagonist activity was to challenge effects of the agonist (R) α -methylhistamine. Therefore, the first objective was to establish a dose response curve for behavioral effects of (R) α -methylhistamine. Male albino CF-1 mice weighing 20-30 g were used. Saline or (R) α -methylhistamine in saline was injected i.p. in a volume \leq 0.4 ml. Animals were observed for various behaviors three times for 10 seconds during each 10 minute interval for a total of 2 hours. Animals were scored for the presence (1) or absence (0) of the behavior and the results were reported as the accumulated score for a 30 minute period (maximum score = 9). As shown in FIG. 4, (R) α -methylhistamine produced a dose-dependent (range of 15 to 35 mg/kg) increase in sleeping one hour after injection. The effect was also evident at 30 minutes after injection.

To assess the effects of antagonists, they were administered with the (R) α -methylhistamine in saline. FIG. 5 shows that thioperamide was able to inhibit the soporific effect of 30 mg/kg of (R) α -methylhistamine. With thioperamide alone (i.e., in the absence of the α -methylhistamine H_3 receptor agonist), animals were

SUBSTITUTE SHEET (RULE 26)

very active, exhibiting normal behaviors. FIG. 6 shows that compound 1 inhibited the soporific effect of 25 mg/kg (R) α -methylhistamine.

5

7.3. DISCUSSION

The results of the *in vitro* (see section 6, *supra*) and *in vivo* activity assays show that a compound of the invention is useful for increasing histamine activity in the brain.

10

In the foregoing *in vivo* assays, thioperamide was used as a positive control. The results indicate that compound 1 is effective as an H₃-receptor antagonist. Direct comparison of the two compounds is not available from the data, however, since the experimental protocols used to test each were not identical.

15

It is noteworthy that in all testing to date, no toxicity of the 1 compound has been observed, even at high doses.

20

8. SPECIFICITY OF COMPOUND 1

The selectivity of action of compound 1 for histamine H₃-receptors was determined in a NOVASCREEN™ receptor selectivity study. At concentrations of 10⁻⁵ M, no significant binding to adenosine, excitatory or inhibitory amino acid, dopamine, serotonin, or a broad range of petidergic receptors, or to ion channel proteins, peptide factor or second messenger systems was observed. The binding study results are shown in Table II.

25

30

35

SUBSTITUTE SHEET (RULE 26)

TABLE II
NOVASCREEN™ RECEPTOR SELECTIVITY ASSAY

5	Receptor/ Selectivity	Reference Compound	Reference K _i (nM)	Initial Percent Inhibition (Average; N=2)
				10 ⁻⁵ M
10	<u>Adenosine</u>			
	Adenosine	NECA	120.00	-3.0
	<u>Amino Acids</u>			
	Excitatory			
	Quisqualate	Quisqualic Acid	11.80	-1.8
15	Kainate	Kainic Acid DME	24.93	42.1
	MK-801	MK801	4.30	-8.6
	NMDA	NMDA	359.00	-4.5
	PCP	PCP	62.30	9.7
	Glycine	Glycine	300.00	1.8
	Inhibitory			
	Glycine	Strychnine Nitrate	33.50	17.4
20	GABA _A	GABA	2.80	0.6
	GABA _B	GABA	176.00	0.0
	Benzodiazepine	Clonazepam	3.40	2.7
	<u>Biogenic Amines</u>			
	Dopamine 1	Butaclamol	37.30	6.4
	Dopamine 2	Spiperone	0.08	3.5
25	Serotonin 1	Serotonin	4.60	-3.6
	Serotonin 2	Serotonin	531.00	10.5
	<u>Peptides</u>			
	Angiotensin	Angiotensin II	0.20	6.5
	Arg-Vasopressin V ₁	arg-Vasopressin	4.90	10.1
	Bombesin	Tyr4-Bombesin	0.55	-5.5
30	CCK Central	CCK	0.13	18.6
	CCK Peripheral	CCK	0.02	6.9
	Substance K	Neurokinin A	2.75	29.2
	Substance P	Substance P	0.08	20.0
	NPY	Neuropeptide Y	0.50	-8.7
	Neurotensin	Neurotensin	1.23	-10.5
35	Somatostatin	Somatostatin	0.03	4.1
	VIP	VIP	1.53	17.1

SUBSTITUTE SHEET (RULE 26)

				Initial Percent Inhibition (Average; N=2)
	Receptor/ Selectivity	Reference Compound	Reference K _i (nM)	10 ⁻⁵ M
	<u>Channel Proteins</u>			
10	Calcium	w-Conotoxin	0.01	1.9
	Calcium	Nifedipine	1.60	8.1
	Chloride	TBPS	112.40	-3.4
	Potassium	Apamin	0.05	7.7
	<u>Peptide Factors</u>			
15	ANF (rat)	ANP	0.15	0.1
	EGF	EGF	0.24	18.1
	NGF	NGF	0.80	17.1
	<u>Second Messenger Systems</u>			
		Forskolin	29.40	2.1
	Adenylate Cyclase Forskolin			
20	Protein Kinase C			
	Phorbol Ester	PDBU	16.50	0.9
	Inositol Triphosphate	IP3	12.50	9.2

25 Values are expressed as the percent inhibition of specific binding and represent the average of duplicate tubes at each of the concentrations tested. Bolded values represent inhibition of fifty percent or greater.

30

35

SUBSTITUTE SHEET (RULE 26)

9. DEMONSTRATION OF APPETITE
SUPPRESSION ACTIVITY IN VIVO

Male Sprague-Dawley rats (200-300 grams) were used in all experiments. Rats were fasted for 24 hours prior to all experiments. Water was provided ad lib. Rats were singly placed in a ventilated plastic circular rat metabolism chamber with a stainless steel wire mesh grid floor equipped with food and H₂O attachments. A measured quantity of food (50 grams) and water (100 mls) was supplied through attachments to the metabolism chambers. Food and water intake was measured at the 2, 4 and 6 hour time periods.

Immediately prior to placement of the rats in the metabolism chambers, animals were administered interperitoneally, vehicle or compound 1 (3, 10 or 30 mg/kg). Final volumes were 1.0 ml/kg. Subsequently, they were left undisturbed for the experimental period except when the food and H₂O attachments were briefly removed for measurement of food and water intake. At the completion of the experiment, animals were returned to their home cages. The results are found in Table III and Figures 7 and 8.

Table III
The Effect of Food Intake In Rats
Following The Administration of Compound I

	Column 1	2 HR	S.E.	4 HR	S.E.	6 HR	S.E.
1	VEHICLE	6.167 ± 0.366		7.083 ± 0.358		13.000	
2	3 mg/kg	5.687 ± 0.494		7.333 ± 0.558			
3	10 mg/kg	2.000 ± 0.258		5.167 ± 0.601			
4	30 mg/kg	1.167 ± 0.477		2.333 ± 0.615		5.000 ± 0.940	

The above protocol is repeated for the histamine H₃ receptor antagonists disclosed herein.

SUBSTITUTE SHEET (RULE 26)

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become
5 apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

10 Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

15

20

25


30

35

SUBSTITUTE SHEET (RULE 26)

WHAT IS CLAIMED IS:

1. Use of a compound of the formula:



wherein R_1 represents hydrogen, an *in vivo* hydrolyzable group, an alkyl group, an cyclic alkyl group, or an aryl group; D is CH_2 or CH_2CH_2 ; Z is S or O; x is 0 or 1; n is an integer from 0 to 6; and R_2 represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 atoms, or a substituted or unsubstituted aryl group of up to about 20 carbon atoms, with the provisos that if R_2 is *tert*-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0; and if R_2 is adamantane, the sum of x and n must be greater than 1;


25 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for suppressing appetite in a subject.

30

35

SUBSTITUTE SHEET (RULE 26)

2. The use according to claim 1 in which said compound is a compound of the formula:



wherein x is 0 or 1; n is an integer from 0 to 6; and R represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 carbon atoms, or substituted or unsubstituted aryl group of up to about 20 carbon atoms, with the provisos that if R is tert-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0; and if R is adamantane, the sum of x and n must be greater than 1;

or a pharmaceutically acceptable salt thereof.


25

30

35


SUBSTITUTE SHEET (RULE 26)

3. The use according to claim 2 in which said compound is a compound of the formula:



in which n is 1, 2, 3 or 4; or a pharmaceutically acceptable salt thereof.

4. The use according to claim 2 in which said compound is a compound of the formula:




25 or a pharmaceutically acceptable salt thereof.

30

35


SUBSTITUTE SHEET (RULE 26)

5. The use according to claim 2 in which said compound is a compound of the formula:



wherein n is 0, 1, 2, 3 or 4; or a pharmaceutically acceptable salt thereof.


6. The use according to claim 2 in which said compound is a compound of the formula:



25 or a pharmaceutically acceptable salt thereof.


30
35
SUBSTITUTE SHEET (RULE 26)

7. The use according to claim 2 in which said compound is a compound of the formula:



in which n is 1, 2, 3 or 4; or a pharmaceutically acceptable salt thereof.

8. The use according to claim 2 in which said compound is a compound of the formula:




or a pharmaceutically acceptable salt thereof.

SUBSTITUTE SHEET (RULE 26)

9. The use of compound 4-(1-cyclohexylvaleroyl-4-piperidyl)1H-imidazole or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for suppressing appetite in a subject.

10. A pharmaceutical composition comprising an amount of a compound of the formula:




wherein R_1 represents hydrogen, an *in vivo* hydrolyzable group, an alkyl group, an cyclic alkyl group, or an aryl group; D is CH_2 or CH_2CH_2 ; Z is S or O; x is 0 or 1; n is an integer from 0 to 6; and R_2 represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 atoms, or a substituted or unsubstituted aryl group of up to about 20 carbon atoms, with the provisos that if R_2 is *tert*-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0; and if R_2 is adamantane, the sum of x and n must be greater than 1;

or a pharmaceutically acceptable salt thereof, effective to achieve appetite suppression, and a pharmaceutically acceptable carrier or excipient.

SUBSTITUTE SHEET (RULE 26)

11. The pharmaceutical composition according to claim 10 wherein said compound is a compound of the formula:

5



10

15 wherein x is 0 or 1; n is an integer from 0 to 6; and R represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 carbon atoms, or substituted or unsubstituted aryl group of up to about 20 carbon atoms, with the provisos that if R is tert-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0; and if R is adamantane, the sum of x and n must be greater than 1;

25


or a pharmaceutically acceptable salt thereof.

30

35

SUBSTITUTE SHEET (RULE 26)

12. The pharmaceutical composition according to claim 11 wherein said compound is a compound of the formula:




15 in which n is 1, 2, 3 or 4; or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical composition comprising an amount of 4-(1-cyclohexylvaleroyl)-4-piperidyl)-1H-imidazole or a pharmaceutically acceptable salt thereof, effective to achieve appetite suppression, and a pharmaceutically acceptable carrier or excipient.

SUBSTITUTE SHEET (RULE 26)

14. A method of suppressing appetite in a subject comprising administering to a subject in whom appetite suppression is desired, an effective amount of a compound of the formula:



15 wherein R_1 represents hydrogen, an *in vivo* hydrolyzable group, an alkyl group, an cyclic alkyl group, or an aryl group; D is CH_2 or CH_2CH_2 ; Z is S or O; x is 0 or 1; n is an integer from 0 to 6; and R_2 represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 atoms, or a substituted or unsubstituted aryl group of up to about 20 carbon atoms, with the provisos that if R_2 is *tert*-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0; and if R_2 is

20

25 adamantane, the sum of x and n must be greater than 1;


or a pharmaceutically acceptable salt thereof.

30

35

SUBSTITUTE SHEET (RULE 26)

15. The method according to claim 14 in which the compound is a compound of the formula:




wherein x is 0 or 1; n is an integer from 0 to 6; and R represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 carbon atoms, or substituted or unsubstituted aryl group of up to about 20 carbon atoms, with the provisos that if R is tert-butyl, cyclohexyl, or dicyclohexylmethyl, x or n must not be 0; and if R is adamantane, the sum of x and n must be greater than 1;

or a pharmaceutically acceptable salt thereof.


SUBSTITUTE SHEET (RULE 26)

16. The method according to claim 15 in which the compound is a compound of the formula:



in which n is 1, 2, 3 or 4; or a pharmaceutically acceptable salt thereof.

17. The method according to claim 15 in which the compound is a compound of the formula:




or a pharmaceutically acceptable salt thereof.

30

35


SUBSTITUTE SHEET (RULE 26)

18. The method according to claim 15 in which the compound is a compound of the formula:



wherein n is 0, 1, 2, 3 or 4; or a pharmaceutically salt thereof.

19. The method according to claim 15 in which the compound is a compound of the formula:




25 or a pharmaceutically acceptable salt thereof.

30

35


SUBSTITUTE SHEET (RULE 26)

20. The method according to claim 15 in which the compound is a compound of the formula:



in which n is 1, 2, 3 or 4; or a pharmaceutically acceptable salt thereof.

15 21. The method according to claim 15 in which the compound is a compound of the formula:



or a pharmaceutically acceptable salt thereof.

30

35

SUBSTITUTE SHEET (RULE 26)

22. The method according to any one of claims 14 to 21 in which the subject is human, and said administration is to treat obesity or weight gain.

5 23. The method according to claim 22 wherein said compound is administered orally.

24. The method according to claim 22 wherein said compound is administered parenterally.

10

25. The method according to claim 22 wherein the amount of said compound administered is about 0.01 mg/kg to about 200 mg/kg in a single dose or divided dose per day.

15

26. A method of suppressing appetite in a subject comprising administering to a subject in whom appetite suppression is desired, an effective amount of 4-(1-cyclohexylvaleroyl-4-piperidyl)1H-imidazole or
20 a pharmaceutically acceptable salt thereof.

27. The method according to claim 26 in which the subject is human, and said administration is to treat obesity or weight gain.

25

28. The method according to claim 27 wherein said compound is administered orally.

29. The method according to claim 27 wherein
30 said compound is administered parenterally.

30. The method according to claim 27 wherein the amount of said compound administered is about 0.01 mg/kg to about 200 mg/kg in a single dose or divided
35 dose per day.

SUBSTITUTE SHEET (RULE 26)

1/8




FIG.1

SUBSTITUTE SHEET (RULE 26)




FIG.2

SUBSTITUTE SHEET (RULE 26)

3/8




FIG.3

SUBSTITUTE SHEET (RULE 26)

4/8




FIG.4

SUBSTITUTE SHEET (RULE 26)

5/8




FIG.5

SUBSTITUTE SHEET (RULE 26)

6/8




FIG.6

SUBSTITUTE SHEET (RULE 26)

7/8





FIG.7

SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/11790

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07D 401/04; A61K 31/445

US CL : 546/203, 210; 548/314.7; 514/319, 326, 397, 910

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/203, 210; 548/314.7; 514/319, 326, 397, 910

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS Online

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, A, 0,197,840 (ARRANG ET AL.) 15 OCTOBER 1986, see entire document.	1-46
A	US, A, 5,051,424 (WIERINGA) 24 SEPTEMBER 1991, see entire document.	1-46
A	US, A, 4,220,653 (VIVINO) 02 SEPTEMBER 1980, see entire document.	1-46

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A		document defining the general state of the art which is not considered to be of particular relevance
* E		earlier document published on or after the international filing date
* L		document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)
* O		document referring to an oral disclosure, use, exhibition or other means
* P		document published prior to the international filing date but later than the priority date claimed
	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is considered with one or more other such documents, such combination being obvious to a person skilled in the art
	* Z	document member of the same patent family

Date of the actual completion of the international search
28 NOVEMBER 1994

Date of mailing of the international search report

13 JAN 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

MICHAEL B. HYDORN

Facsimile No. (703) 305-3230

Telephone No. (703) 305-1235

Form PCT/ISA/210 (second sheet)(July 1992)*